



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

a sustained flight of approximately forty miles. STEPHEN S. GREGORY JR., 456 Surf St., Chicago, Ill.

A Crowd of Hummers.—On May 1, 1920, after I had returned from a remarkable field trip, on which I had found 94 species of birds, one of my neighbors, Colonel B. F. Procter, called me to see the Ruby-throated Hummingbirds about a red-flowered horse-chestnut on his lawn. There was a continuous going and coming of the hummers, twenty or more being in sight all the time. I saw several perched on the smaller twigs of the tree, in addition to the group buzzing about the fragrant blossoms. Though these tiny birds came every day for a week, I never again saw so many at one time. I feel sure that I saw on that one day more individuals of this species than I have ever seen in any other whole season. This bird is usually so rare that I keep a record of every individual seen.—GORDON WILSON, Bowling Green, Kentucky.

Plumage of the Chimney Swift.—Dr. Wetmore's interesting article on the wing claws of *Chaetura pelagica*, in the December 'Condor,' reminded me of another curious fact regarding this bird.

In May, 1898, I put up a lot of skins of Chimney Swifts which had been suffocated in a chimney at Waukegan, Ill., and found concealed among the feathers of the abdomen at about the middle, a tuft of pure white down. This could only be noticed on parting the feathers, and was present in all of them but was lost or absorbed in making up the skins. Last Spring, while preparing a specimen (with the wings spread) my attention was called to a tract of naked black skin on the underside of the wing joint, and upon closer inspection, I found a single smoky black oval shaped feather growing from the center of the naked skin.

Upon examining a number of other specimens of *Chaetura pelagica*, and of *C. vauxi* from California and *C. caudacuta* from New South Wales and Japan, the single feather was also found.—HENRY K. COALE, Highland Park, Ill.

Mortality among Chimney Swifts.—In 'The Auk', Vol. XXV, No. 3, pp. 317-318, July, 1908, under the heading "Curious fatality among Chimney Swifts," Mr. Ruthven Deane has recorded that some 700 Chimney Swifts (*Chaetura pelagica*) entered a chimney in the house of Dr. Maxson, Waukegan, Ill., during a heavy rainstorm, while gases from a furnace fire were passing off through the chimney. More than 100 of these birds were killed, and the rest were released while yet alive by the occupants of the house. In another chimney of the same house no Swifts were found at that time.

A destruction of Chimney Swifts in even greater numbers, at Truro, Nova Scotia, under circumstances similar, yet different, is described in the following extract from a letter dated June 1, 1919, addressed to me by Mr. E. C. Allen, now of Halifax, Nova Scotia, to whom I am indebted for permission to publish this note.